Site: SORGHUM	Overall Confidence Rating: M								
Background: Total acres planted was approximately 13, 470,000 <sup>1</sup> .									
Organophosphate	% Tr	eated	# Applications/yr		Rate (lb A	I/A)/yr	PHI (days)		
Pesticides	Max <sup>5</sup>	Avg <sup>5</sup>	Max	Avg	Max	Avg <sup>5</sup>	Min	Avg	
chlorpyrifos	5	3	2 3	1 5	1 <sup>6</sup>	0.5	30 <sup>3</sup> 60 <sup>6</sup>		
diazinon <sup>D</sup>					4 6		7 6		
dimethoate	1	1	3 <sup>3</sup>		0.5 4		28 3		
disulfoton	1	<1	3 <sup>3</sup>	1 5	1 4	0.5	7 <sup>3</sup> 60 <sup>4</sup>		
malathion	0.3	0.1	as needed	1.0 5	0.95 6	1.1	37 <sup>6</sup>		
methamidophos	<1	<1		1.0 5		1			
methidathion			3 6		0.5 6		30 <sup>6</sup>		
oxydemeton-methyl	<1	<1		1.0 5		0.24	_		
parathion, ethyl		1			0.95 6		1221 <sup>6</sup>		
phorate	1	1	2 4	1 <sup>3</sup>	1 4		28 6		
terbufos	12		1.0 <sup>3</sup>		3.92 <sup>6</sup>				

note: Acephate does not have a tolerance listed for use on sorghum (see Tolerance Index System), as of 7/2/98.

Confidence Rating: H= high confidence = data from several confirming sources; confirmed by personal experience

M = medium confidence = data from only a few sources; may be some conflicting or unconfirmed info.

L = low confidence = data from only one unconfirmed source

Organophospha	ate Target Pests for (YOUR CROP) (Primary pests controlled by the OP's)
Major	aphids, wireworms, cutworms
Moderate	stink bugs, chinch bugs, rootworms
Minor	grasshoppers, ants, other lepidopteran larvae, weevils, beetle grubs, midges, borers, fleahoppers, mites, thrips, seed-corn beetles, false chinch bugs, nematodes, sorghum webworm.

Major = 20+% of all OP usage on pest; Moderate = 5-20% of all OP usage on pest; Minor =<5% of all OP usage on pest

## **Sources:**

- (1). Doane 1996
- (2). QUA Plus
- (3). State Recommendations
- (4). Insect Control Guide 1997 (Vol. 9)
- (5). QUA
- (6). LUIS

## **Key to Timing** (see Pest Summary Matrix)

A = seed timing

B = at plant timing

C = postemergence/foliar spray timing

D = boot timing

Date: 6/15/98

<sup>\*</sup> Superscript numbers refer to resources used; superscript letters refer to timing of applications (see last page).

Pest	Organophosphate	Efficac y	Mkt	Class	Alt. Pesticide List	Efficacy	Mkt	Comments on Alternatives
Timing: see key to timing								
Aphids	chlorpyrifos	O 2	med.	С	aldicarb		low	
[greenbugs, corn leaf aphids, etc.]	dimethoate		low	С	carbofuran	⊚ <sup>2</sup>	low	
(major)	disulfoton <sup>B</sup>		med.	P	lambda-cyhalothrin		low	
aphidids	malathion <sup>2</sup>			P	permethrin		low	
	parathion, ethyl		med.	С	thiodicarb		low	
	phorate <sup>B</sup>		low					
	oxydemeton-methyl <sup>3</sup>							
	terbufos 3 B	O 2	high					
Wireworms	chlorpyrifos		high	С	carbofuran <sup>3</sup>	O 3		
(major)	diazinon <sup>3 A</sup>			О	imidacloprid <sup>3 A</sup>			
elaterids	terbufos 3 B		high	СН	lindane <sup>3 A</sup>			
				P	permethrin		med.	
Cutworms	chlorpyrifos	O 3	high	Р	esfenvalerate		low	
[black cutworm] (major)	terbufos <sup>3 B</sup>		med.	Р	lambda-cyhalothrin		low	
noctuids				P	permethrin		med.	

Efficacy Rating: Excellent = ⊚ Good = O Fair = ●

Market Share: High = use of OP represents 20+% of all insecticide usage on pest; Med = 5-20% of all usage on pest; Lo = <5% of all usage on pest Insecticides: C = Carbamates; P = Pyrethroids; CH = Chlorinated Hydrocarbons; IGR = Insect Growth Regulators; B = Biological; O = Other pesticides

<sup>\*</sup> Superscript numbers refer to resources used; superscript letters refer to timing of application (see last page for details).

Pest Importance: Major = 20+% of all OP usage on pest; Moderate = 5-20% of all OP usage on pest; Minor = <5% of all OP usage on pest

Pest	Organophosphate	Efficac y	Mkt	Class	Alt. Pesticide List	Efficacy	Mkt	Comments on Alternatives
Timing: see key	Timing: see key to timing							
Hemipteran	chlorpyrifos 3 C		med.	С	aldicarb <sup>3 B</sup>			note: cyfluthrin – is a restricted use chemical.
Bugs [chinch bugs,	diazinon <sup>3 A</sup>			С	carbaryl <sup>3 C</sup>		low	note: carbofuran - use precautions.
green stink bugs] (moderate)	dimethoate		low	С	carbofuran <sup>3 B, 3 C</sup>		high	
	disulfoton		med.	P	cyfluthrin <sup>3</sup>			
lygaeids pentatomids	parathion, ethyl		low	P	esfenvalerate		low	
	phorate		low	СН	endosulfan		low	
	terbufos <sup>3 B</sup>		high	О	imidacloprid <sup>3 A</sup>			
				СН	lindane <sup>3 A</sup>			
				С	methomyl		low	
				С	thiodicarb		low	
Rootworms	terbufos		high	С	carbofuran		med.	
(moderate)				Р	lambda-cyhalothrin		med.	
chrysomelids				Р	tefluthrin		low	
Grasshoppers	chlorpyrifos		low	С	carbaryl		low	
(minor)	dimethoate		med.	P	lambda-cyhalothrin		med.	
acridids	parathion, ethyl		high					

Pest Importance: Major = 20+% of all OP usage on pest; Moderate = 5-20% of all OP usage on pest; Minor = <5% of all OP usage on pest

Efficacy Rating: Excellent =  $\bigcirc$  Good =  $\bigcirc$  Fair =  $\bigcirc$ 

Market Share: High = use of OP represents 20+% of all insecticide usage on pest; Med = 5-20% of all usage on pest; Lo = <5% of all usage on pest Insecticides: C = Carbamates; P = Pyrethroids; CH = Chlorinated Hydrocarbons; IGR = Insect Growth Regulators; B = Biological; O = Other pesticides

<sup>\*</sup> Superscript numbers refer to resources used; superscript letters refer to timing of application (see last page for details).

Pest	Organophosphate	Efficac y	Mkt	Class	Alt. Pesticide List	Efficacy	Mkt	Comments on Alternatives
Timing: see key to timing								
Ants (minor)	chlorpyrifos		low	О	cryolite		med.	
formicids	terbufos <sup>3 B</sup>		high					
Lepidopterous	chlorpyrifos		med.	В	Bacillus thuringiensis	•	low	
Larvae - others (minor)	dimethoate		low	Р	cyfluthrin		med.	
[corn earworm,	parathion, ethyl		low	P	cypermethrin		low	
fall armyworm] noctuids				СН	endosulfan		low	
noctures				P	esfenvalerate		low	
				P	lambda-cyhalothrin		high	
				С	methomyl		med.	
				P	permethrin		low	
				С	thiodicarb		low	
				P	zetamethrin		low	
Midges	chlorpyrifos		low	О	barium polysulfide		low	
(minor) [sorghum midge] cecidomyiids	dimethoate		low	P	esfenvalerate		low	
				P	lambda-cyhalothrin		high	
				P	permethrin		low	
				Р	zetamethrin		low	

<sup>\*</sup> Superscript numbers refer to resources used; superscript letters refer to timing of application (see last page for details).

Pest Importance: Major = 20+% of all OP usage on pest; Moderate = 5-20% of all OP usage on pest; Minor = <5% of all OP usage on pest

Efficacy Rating: Excellent = ⊕ Good = O Fair = ●

Market Share: High = use of OP represents 20+% of all insecticide usage on pest; Med = 5-20% of all usage on pest; Lo = <5% of all usage on pest Insecticides: C = Carbamates; P = Pyrethroids; CH = Chlorinated Hydrocarbons; IGR = Insect Growth Regulators; B = Biological; O = Other pesticides

## ADDITIONAL INFORMATION:

Keys to Timing: (A) seed timing; (B) at plant timing; (C) postemergence/foliar timing spray; (D) boot timing.

## **RESOURCES:**

Keys to Resources:

- (1). Doane 1996
- (2). QUA plus (database and recent arrivals)
- (3). State Recommendations
- (4). Insect Control Guide, Vol. 9 (1997)
- (5). QUA
- (6). LUIS

Date: 6/12/98